

Progress in ERAWAST (Exotic Radionuclides from Accelerator Waste for Science and Technology)

Dorothea Schumann
Paul Scherrer Institute
RadWasteAnalytics

Outline

- Objective of the project
- ERAWAST plan
- Isotope production sources
 - ◆ Copper beam dump
 - ◆ Lead targets from SINQ
- Summary and Outlook

Objective

Application of exotic long-lived isotopes from accelerator waste for scientific purposes

Examples:

^7Be (50.8d), ^{10}Be ($1.6 \cdot 10^6\text{y}$), ^{26}Al ($7.2 \cdot 10^5\text{y}$), ^{44}Ti (60.4y), ^{53}Mn ($3.7 \cdot 10^6\text{y}$), ^{60}Fe ($1.5 \cdot 10^6\text{y}$), ^{146}Sm ($1.03 \cdot 10^8\text{y}$), ^{182}Hf ($9 \cdot 10^6\text{y}$), ^{129}I ($1.56 \cdot 10^7\text{y}$)

Development of a long-term international collaboration between

- ◆ Nuclide production facilities
- ◆ Basic physics research
- ◆ Nuclear Data for P&T
- ◆ Nuclear astrophysics
- ◆ AMS measurement groups
- ◆ Pharmaceutical chemistry
- ◆ Nanotechnology

ERAWAST – plan

1. Existing accelerator waste material

Copper beam dump irradiated at the 590-MeV proton beam station at PSI, dismantled about 15 years ago

^{26}Al , ^{59}Ni , ^{53}Mn , ^{60}Fe , ^{44}Ti or others can be separated

other irradiated materials like carbon (^{10}Be), stainless steel or concrete are also available

1 PostDoc position financed from PSI for the development of separation in a technological scale now available

2. Target material from the SINQ facility

Two irradiated lead targets from the spallation source are available.

Heavier isotopes like ^{182}Hf , ^{129}I or several rare earth elements (e.g.

^{146}Sm , several Dy isotopes) can be obtained. In principle, targets from the SINQ will be available every second year.


3. Special irradiations

The SINQ facility offers the possibility to irradiate materials with 590 MeV protons at special positions. Tended experiments for isotope production can be offered.

Characteristics of the copper beam dump


- Beam stop from the former BMA station
- Operated from 1980-1992, dismantled in 1993
- 0.1 Ah total beam dose (590 MeV protons)
- copper cylinder of ~ 10 kg; diameter 80mm
- Sample taking from several parts by drilling

Schematic view of the beam dum



* - Area of drilling $\text{Ø } 20\text{mm}$



Before drilling



Sample	⁴⁴ Ti [kBq/g]	³⁶ Cl [Bq/g]	⁶³ Ni [kBq/g]	⁵⁵ Fe [kBq/g]	⁶⁰ Fe [Bq/g]	²⁶ Al [Bq/g]	^{110m} Ag [Bq/g]	^{108m} Ag [Bq/g]	⁵⁹ Ni [Bq/g]	⁵³ Mn [Bq/g]	⁶⁰ Co [kBq/g]
C6.1.1	4.8	4.06	220.3	233.6		0.154	1.21	1.77	594		224.0
C6.1.2	0.54	0.49	133.7	37.3		0.016	1.98	2.52			85.7
C6.2.1	1616.0		34151.1	42450.3	84.7	140	3.89	62.58	127000	6900	49957.1
C6.2.2	0.2	0.30	217.7	108.6		0.01	2.48	1.35	129	0.6	111.6
C6.3.1	740.8		36566.7	44136.3	58.5	56	2.3	37.27	125000	4310	37969.6
C6.3.2	18.6	0.19	2006.0	2562.0	3.3	1	0.453	9.19	6620	112	2691.8
C6.3.3	1.5	0.02	1109.7	1552.4	1.9	0.2	1.32	9.22	2620	117	1239.4
C6.3.4	0.6	0.01	1841.6	257.0	0.5	0.03	0.92	1.69	759	6	663.2
C6.3.5	0.4	0.35	706.2	154.7	0.5	0.01	0.56	1.06	466	4.9	438.8
C6.4.1	778.1		16776.3	26590.4	52.2	41	1.64	27.70		3600	47256.0
C6.4.2		0.24	799.1	132.5		0.008	0.09	1.27		2.1	505.5
C6.5.1	95.0		5764.1	11520.4	20.2	3	11.70	13.64	25700	998	10091.9
C6.5.2		0.27	545.6	157.8		0.01	0.59	1.80	422	2.0	415.0
C6.6.1	-	0.13	1005.7	287.6	0.7	0.012	3.75	3.93	6940	10.4	459.0
C6.6.2	-	0.08	233.2	127.8		0.0019	0.49	0.94		0.6	169.8
C6.7.1	-	0.08	170.2	350.7	0.7	0.005	1.4	1.34		1.4	148.6
C6.7.2	-	0.04	118.8	233.6		0.0013	1.85	0.86		0.5	91.1
C6.7.3	-	0.04			0.1	0.0012	3.86	1.72	1		56.1
C6.7.4	-	0.04				0.0009	0.18	1.03		0.5	6.1

Nuclide distribution
dependent on nuclear
reaction and particle
energy

rel. frequency [%]



Estimation of available radionuclides (no separation)

Basis: central drilling with 2 cm diameter

500g copper

Average of all 4 central values

Average of the central value and from 1 cm distance

„Safety factor“ of 2

^{44}Ti : 100 MBq (10^{18} atoms)

^{53}Mn : 500 kBq (10^{19} atoms)

^{26}Al : 7 kBq (10^{17} atoms)

^{60}Fe : 5 kBq (10^{17} atoms)

^{59}Ni : 8 MBq (10^{19} atoms)

(^{60}Co : 5 GBq)

All these radionuclides can be provided without carrier, but some of them contain other long-lived isotopes ($^{55}\text{Fe}/^{63}\text{Ni}$)

Preparation of ^{60}Fe -samples

- ◆ Chemical separation of the iron fraction from samples of the copper beam-dump (590 MeV protons, beam dosis ca. 0.1Ah); $\sim 10^{15}$ - 10^{16} atoms
- ◆ Basic research:
 - ◆ Determination of the **half-life of ^{60}Fe** (collaboration with TU Munich, Germany)
 - ◆ Cross section measurement of the reaction **$^{60}\text{Fe}(n,\gamma)^{61}\text{Fe}$** (collaboration with FZK, Germany)
- ◆ Applied research: Preparation of standard solutions for accelerator mass spectrometry, biomedical investigations, astrophysics

Chemical separation of ^{60}Fe

Special Problem:



^{60}Fe : no γ radiation, low β -energy

Measurement of the increase of the Co-daughter




Measurement of the ^{61}Fe production (1027/1205 keV)

→ very good chemical separation from Co necessary

- Dissolution of Cu chips (beam dump) in 7 M HNO_3
- Evaporation to dryness
- Dissolution in 7 M HCl
- + 5 mg Fe^{3+} and 5 mg Co^{2+} as carrier
- Extraction with diisopropylether
- Aqueous phase: Ni, Co, Cu, organic phase: Fe
- Back extraction with 0.1 M HCl , repetition of procedure
- Additional purification by precipitation of $\text{Fe}(\text{OH})_3$
- Result: $3.8 \cdot 10^{15}$ (TUM) / $1.2 \cdot 10^{16}$ (FZK) ^{60}Fe atoms, decontamination factor (Co) $> 10^8$ (0.3 Bq)

Lead targets from SINQ

2 Samples from target 4, 2 years operation; EOB 1999



Analysis results

	D9 [Bq/g]	D14 [Bq/g]		D9 [Bq/g]	D14 [Bq/g]
^{207}Bi	$3.00 \cdot 10^7$	$1.01 \cdot 10^7$	^{106}Ru	$4.83 \cdot 10^6$	$3.91 \cdot 10^6$
$^{172}\text{Lu}/^{172}\text{Hf}$	$2.00 \cdot 10^7$	$5.41 \cdot 10^7$	$^{110\text{m}}\text{Ag}$	$1.29 \cdot 10^6$	$3.93 \cdot 10^5$
^{173}Lu	$2.76 \cdot 10^7$	$4.30 \cdot 10^7$	^{125}Sb	$1.32 \cdot 10^6$	-
$^{194}\text{Au}/^{194}\text{Hg}$	$1.86 \cdot 10^7$	$3.13 \cdot 10^6$	^{133}Ba	$2.8 \cdot 10^6$	$7.94 \cdot 10^5$
^{102}Rh	$5.53 \cdot 10^6$	$1.44 \cdot 10^5$	$^{44}\text{Sc}/^{44}\text{Ti}$	$8.00 \cdot 10^4$	$2.84 \cdot 10^4$
$^{202}\text{Tl}/^{202}\text{Pb}$	$4.80 \cdot 10^5$	$1.87 \cdot 10^5$	$^{108\text{m}}\text{Ag}$	$3.75 \cdot 10^5$	$1.56 \cdot 10^4$
^{60}Co	$3.67 \cdot 10^6$	$1.40 \cdot 10^6$	$^{194}\text{Os}/^{194\text{m}}\text{Ir}$	$2.61 \cdot 10^4$	-
^{54}Mn	$2.29 \cdot 10^5$	$7.01 \cdot 10^4$	^{26}Al	0.5	0.2
^{58}Co	$1.55 \cdot 10^6$	$9.47 \cdot 10^5$	^{36}Cl	$9.5 \cdot 10^1$	$4.8 \cdot 10^1$
^{55}Fe	$8.73 \cdot 10^7$	$5.99 \cdot 10^7$	^{63}Ni	$6.30 \cdot 10^8$	$4.52 \cdot 10^8$

Summary

- ERAWAST-community established (appr. 30 partners, growing number), new proposal for Network Programme
- First samples provided: ^{10}Be (GSI), ^{60}Fe (TUM; FZK), ^{26}Al , ^{44}Ti (Uni Mainz; Edinburgh)
- Up to 10^{18} atoms of ^{10}Be available (separated)
- 10^{16-19} atoms of several radionuclides (^{26}Al , ^{60}Fe , ^{59}Ni , ^{53}Mn , ^{44}Ti) available (not yet separated)
- PostDoc for subsequent separation from Cu in a Hot-cell available
- Analytical work on Pb-targets ongoing, heavier isotopes can in principle be provided
- More Cu-, Pb- and C-samples available
- Possibilities for other irradiation positions (SINQ, beam dumps, collimators)
- Problem of mass separation still unsolved